

STUDY THE EFFECT OF SUBSTRATE AND NON SUBSTRATE MEDIUM ON THE PECTINASE ACTIVITY OF FUNGI FROM VEGETABLES WASTE

*Balwan W. Kamble, *Shrimant A. Survase,*Anil U. Kulkarniand**Ashok M. Chavan

*Department of Botany, Lalbahadur Shastri Sr. College, Partur Dist. Jalna.431501.

**Department of Botany, Dr. Babasaheb Ambedkar Marathwada University, Aurangabad – 431004 (M.S.) India

Email-kamblebalwan@gmail.com

Abstract


Many plants cell wall contain pectin and it degrades by the microorganisms for their growth. Pectinase have been used in several conventional industrial processes over the years, such as textile, plant fiber processing and tea. In order to study the production of pectinase by ten dominating fungal species associated with vegetable waste were tested on substrate (pectin nitrate) and non substrate (glucose nitrate) medium. Pectinase activity was determined by viscometrical method.

The result revealed productions of pectinase were tested on substrate and non substrate broth medium. The fungi were subjected to different time intervals and pectinase enzymatic activity representing percent loss of viscosity. The highest pectinase activity was obtained by *Penicillium notatum*, *Trichoderma viride* and *Aspergillus niger* as compare to other fungi like *Aspergillus flavus*, *A. fumigatus*, *Alternaria alternata*, *Curvularia lunata*, *Fusarium oxysporum*, *Penicillium spp.* and *Rhizopus stolonifer* at different interval time 20, 40, 60 min.

Key word: Substrate, non substrate medium, Ostwald viscometer, fungi, pectinase activity, vegetable waste.

Introduction

The vegetables get spoiled by several biotic problems and an under threat due to fungi in all over at vegetables at market areas. Soil-borne plant fungi cause significant damage to almost all crops particularly to the vegetables (Usman *et al.*, 2013). In India, 20-30 % of the produce is spoiled in the markets (FAO, 2002; Deka *et al.*, 2006). A huge amount of these materials to be decomposed by microorganism such as bacteria and fungi. Pectinase enzyme are produced by microorganisms including fungi, yeasts and bacteria, (Cao *et al.*, 1992; Blanco *et al.*, 1999; Huang and Mahoney,



1999). Pectinase, include a group of enzymes that are responsible for the degradation of pectin substances and have important applications in the food industry (Sandri, I.G. et.al, 2011)& (Sandri, I.G.et.al, 2013).

Now a day, pectinase enzyme is one of the most important enzymes in food processing industries mainly for extraction and clarification of fruit juices and wines (Oyewole et al., 2011). Pectinase have been used in several conventional industrial processes over the years, such as textile, plant fiber processing, tea and coffee industries, oil extraction, treatment of industrial waste water, containing pectinacious material, and paper manufacturing (Jayani et al, 2005).

Application of agro-industrial residues as carbon sources in enzyme production processes reduces the cost of production, and also helps in solving problems with their disposal (Murad and Azzaz, 2010).Such residues have yielded good results in the production of pectinase (Silvaet al., 2002). Pectinase constitute approximately 10% of the total enzyme production in the world market and 25% of global sale in the food industry, Naderi, S. et.al (2012).


In present investigation attempts were made to study the pectinase enzymes activity by dominating fungi from vegetable waste.

Material and Methods

Production of pectinase

Production of pectinase was made by growing the fungi in liquid medium containing pectin – 10gm, KNO_3 – 0.25%, KH_2PO_4 – 0.1%, $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ – 0.05%, pH – 5.0. Out of which 25 ml of medium was poured in 100 ml Erlenmeyer conical flasks and autoclaved at 15 lbs pressure for 20 minutes. The flasks on cooling were inoculated separately with 1 ml standard spores suspension of test fungi prepared from 7 days old cultures grown on PDA slants. The flasks were incubated for 6 days at 25⁰C with diurnal periodicity of light. On 7th day, the flasks were harvested by filtering the contents through Whatman filter paper no.1. The filtrates were collected in the presterilized bottles and termed as crude enzyme.

Assay for pectinase



Pectinase activity was assayed by viscometric method (Papdiwal et. al, 1982) as viscosity loss % after 60 minutes. The Ostwald's viscometer was thoroughly cleaned with distilled water and dried before use. 6ml of pectin in 2ml of 0.2 M acetate buffer (pH 5.2) and 4ml of enzyme source were taken in viscometer and were thoroughly mixed and incubated at 25⁰C temperature. The efflux time of the mixture at 0, 20, 40 and 60 minutes was recorded with the help of stop watch.

The percent loss of viscosity was calculated by using the formula,

$$\text{Percent loss of viscosity} = \frac{T_0 - T_x}{T_0 - T_w} \times 100$$

Where, T_0 = Flow time in seconds at zero time

T_x = Flow time of the reaction mixture at time T

T_w = Flow time of distilled water

Biostatic analysis

All the results were statistically analyzed using analysis of variance (ANOVA) test and treatment means were compared using the least significant difference (C.D., $p = 0.05$) which allowed determination of significance between different applications (Mungikar, 1997).

Results and Discussion

In present investigation production of Pectinase by ten dominating fungal species associated with Vegetable waste were tested on substrate (pectin nitrate) and non substrate (glucose nitrate) medium. On the contrary of this pectinase activity of *Aspergillus niger*, *A. flavus*, *A. fumigatus*, *Alternaria alternata*, *Curvularia lunata*, *Fusarium oxysporum*, *Penicillium notatum*, *P.spp*, *Rhizopus stolanifer* and *Trichoderma viride* was stimulated due to pectin.

Photoplate : Fungi cultured on liquid medium



Table1: Effect of non-substrate media on production of pectinase

| Fungi | Non-substrate medium % (Glucose) viscosity loss after min | | | Standard Error (SE) | Critical Difference (CD) p= 0.05 |
|------------------------------|---|-------|-------|------------------------|--|
| | 20 | 40 | 60 | | |
| <i>Aspergillus niger</i> | 41.93 | 44.83 | 46.12 | 1.24 | 5.33 |
| <i>Aspergillus flavus</i> | 03.37 | 04.77 | 05.76 | 0.69 | 2.97 |
| <i>Aspergillus fumigatus</i> | 03.93 | 07.12 | 08.59 | 1.38 | 5.93 |
| <i>Alternaria alternata</i> | 22.29 | 23.99 | 35.66 | 4.20 | 18.06 |
| <i>Curvularia lunata</i> | 20.14 | 22.30 | 25.41 | 1.53 | 6.58 |
| <i>Fusarium oxysporum</i> | 17.82 | 22.48 | 28.29 | 3.03 | 13.03 |
| <i>Penicillium notatum</i> | 03.94 | 07.17 | 09.35 | 1.57 | 6.75 |
| <i>Penicillium spp.</i> | 12.63 | 15.25 | 17.64 | 1.45 | 6.24 |
| <i>Rhizopus stolanifer</i> | 11.10 | 14.32 | 17.90 | 1.96 | 8.43 |
| <i>Trichoderma viride</i> | 02.21 | 04.42 | 05.65 | 1.01 | 4.34 |

Graph 1: Effect of non-substrate media on production of pectinase





Table 2: Effect of substrate media on production of pectinase

| Fungi | Substrate medium % (Pectin) viscosity loss after min | | | Standard Error (SE) | Critical Difference (CD) p= 0.05 |
|------------------------------|--|-------|-------|---------------------|----------------------------------|
| | 20 | 40 | 60 | | |
| <i>Aspergillus niger</i> | 19.67 | 41.16 | 43.77 | 7.64 | 32.85 |
| <i>Aspergillus flavus</i> | 19.39 | 37.79 | 40.13 | 6.56 | 28.21 |
| <i>Aspergillus fumigatus</i> | 01.80 | 04.38 | 07.21 | 1.56 | 6.71 |
| <i>Alternaria alternata</i> | 20.85 | 27.13 | 32.16 | 3.27 | 14.06 |
| <i>Curvularia lunata</i> | 11.11 | 21.11 | 23.33 | 3.76 | 16.17 |
| <i>Fusarium oxysporum</i> | 22.80 | 29.07 | 33.33 | 3.06 | 13.16 |
| <i>Penicillium notatum</i> | 60.37 | 69.81 | 84.90 | 7.14 | 30.70 |
| <i>Penicillium spp.</i> | 12.05 | 23.07 | 31.73 | 5.69 | 24.47 |
| <i>Rhizopus stolanifer</i> | 14.22 | 17.36 | 23.43 | 2.70 | 11.61 |


| | | | | | |
|---------------------------|-------|-------|-------|-------------|--------------|
| <i>Trichoderma viride</i> | 51.72 | 65.51 | 77.58 | 7.47 | 32.12 |
|---------------------------|-------|-------|-------|-------------|--------------|

Graph 2: Effect of substrate media on production of pectinase



The production of pectinase was determined by measuring percent loss of viscosity by the mixture of culture filtrate of substrate and non substrate broth media. The observed records were noted in the table. The percent loss of viscosity was obtained as a particular time intervals. It was cleared from the results table & graph no.1 & 2, summarized that loss of viscosity was directly related with the time. As time interval increased more viscosity loss was observed.

From the result Maximum pectinase activity was detected in non- substrate broth medium from *Aspergillus niger* (46.12% at 60 min.) and minimum (43.77% at 60 min) was showed in substrate medium. Final reading of viscosity loss was increased at time interval. Maximum pectinase activity was detected from *Aspergillus flavus*(40.13 %) in substrate medium and resistant (05.76 % at 60 min) in non-substrate medium. *Aspergillus fumigatus* (08.59% at 60 min), also showed highest pectinase activity in non substrate medium and minimum (07.21% at 60 min)



in substrate medium. The extracellular pectinase activity by *Alternaria alternata* (35.66% at 60 min) in non substrate and (32.16% at 60 min) were exhibited at similar rate. However, *Curvularia lunata* (25.41% at 60 min) showed maximum pectinase activity in non substrate medium, than others (23.33% at 60 min). *Fusarium oxysporum* (33.33% at 60 min) was highest pectinase activity recorded in substrate medium and lowest (28.29% at 60 min) was showed in non substrate medium. The highest pectinase activity obtained from substrate medium by *Penicillium notatum* (84.90 % at 60 min) and lowest found in non- substrate medium. Pectinase activity by *Penicillium spp.* (31.73% at 60 min) also increased in substrate medium and decreased (17.64% at 60 min) in non substrate medium. Maximal pectinolytic activity shown by *Rhizopus stolonifer* (23.43% at 60 min) in substrate medium and minimum in (17.90% at 60 min) non substrate medium. The maximum Pectinase activity detected by *Trichoderma viride* (77.58% at 60 min) in non substrate medium and minimum was showed in substrate medium.


Pectinase activity representing percent loss of viscosity and as time intervals were increased viscosity loss was increased progressively. Similar results were reported by Rathod, 2011, (Bhale and J. N. Rajkonda 2012).

It can concluded from table and graph, that pectinolytic enzyme production was found to be maximum in both substrate and non substrate broth media for all tested fungi in general. *Trichoderma viride* and *Penicillium notatum* showed highest pectinase production on non substrate medium compared to the other fungi. Substrate medium also respond to high pectinase production by *Aspergillus niger* and *Alternaria alternata*. This could be highly beneficial for the production of pectinase from Vegetables waste by fungi.

Acknowledgement

Authors are thankful to Principal of Lal Bahadur Shastri Senior College, Partur for providing all necessary facilities.

Reference:



Blanco, P., C. Sieiro and T.G. Villa (1999). Production of pectic enzymes in yeast. *FEMS Microbiology Letters* 175, 1–9.

Cao, J.; Zheng, L. and S. Chen (1992). Screening of pectinase producer from alkalophilic bacteria and study on its potential application in degumming of ramie. *Enzyme and Microbial Technology*, 14:1013–1016.

Deka, B.C., Choudry, S., Bhattacharya, A., Begum, K.H. and Neog, M. (2006). Post- harvest treatments for shelf life extension of banana under different storage environments. IV International Conference on managing in chains. The integrated view on fruits and vegetables quality. *ISHS ActaHorticulturae*, 712: 110-116.

FAO. (2002). Production year Book, Vol. 54, FAO, Stat. Sec. 163, Food and Agric. Organization, united Nations, Rome.

Huang, L.K. and R.R. Mahoney (1999). Purification and characterization of an endo-polygalacturonase from *Verticillum albo-atrum*. *Journal of Applied Microbiology*, 86:145–146.


Jayani RS , Saxena S & Gupta R. (2005). Microbial pectinolytic enzymes: Areview. *Process Biochemistry Elsevier Ltd*, 40, 2931-2944.

Mungikar, A.M. (1997) “An introduction to Biometry”, Saraswati Printing Press, Aurangabad.

Murad, H.A. and H.H. Azzaz (2010). Cellulase and dairy animal feeding. *Biotechnology*, 9 (3):238-256.

Naderi, S.; Naghavi, N.S.; Shahanipoor, K. (2012)Pectinolytic Activity of *Aspergillus Niger* on Pectic Agricultural Substrates. The 1st International and the 4th National Congress on Recycling of Organic Waste in Agriculture, Isfahan, Iran, Apr 26–27; pp. 1–6.

Oyewole, O.A., Oyeleke, S.B., Dauda, B.E.N. and Emiade, S. (2011). Production of amylase and protease enzymes by *Aspergillus niger* and *Penicillium frequenstans* isolated from abattoir effluent. *Microbiology Journal* 1(5):174-180.



Papdiwal, P. B.(1982) Pectolytic enzymes In: Methods in experimental plant pathology (eds) Mukadam, D. S. and Gangawane, L. V. Marathwada University press Aurangabad pp.; p.14-18.15.

Rathod, G.M. (2011). Studies on post-harvest diseases of papaya. Ph. D. thesis, Dr. Babasaheb Ambedkar Marathwada, University, Aurangabad. pp. 233.

Sandri, I.G.; Fontana, R.C.; Barfknecht, D.M.; Silveira, M.M.(2011) Clarification of fruit juices by fungal pectinases. LWT Food Sci. Technol., 44, 2217–2222.

Sandri, I.G.; Lorenzoni, C.M.T.; Fontana, R.C.; Silveira, M.M.(2013) Use of pectinases produced by a new strain of *Aspergillus niger* for the enzymatic treatment of apple and blueberry juice. LWT Food Sci. Technol, 51, 469–475.

Silva, D.; Martins, E. S.; Silva, R. and Gomes, E (2002). Pectinase production from *Penicillium viridicatum* RFC3 by solid state fermentation using agricultural residues and agro-industrial by-product. *Braz. J. Microbiol.*, **33**, 318-324.

Usman, F., Shaukat, S.S., Abid, M., Hussain, F. (2013). Rhizosphere fungi of different vegetables and their antagonistic activity against pathogenic fungi of brinjal and spinach. *International Journal of Biology and Biotechnology*, **10**: 255-259.

U. N. Bhale and J. N. Rajkonda (2012) Enzymatic activity of *Trichoderma* species. *Novus Natural Science Research, Research*, 1(4), pp. 1-8.